

Emergency Seizure Treatment Step-by-Step

Step 1. Confirm seizure

Signs and Symptoms

When I am having a seizure, I might display some of the following signs or symptoms:

- Convulsions Stiffening
 Unconsciousness Staring
 Involuntary rhythmic movements
 Other _____

Step 2. Provide basic first aid

To ensure my safety, here are some steps to follow:




1. Cushion head, remove glasses.



2. Loosen tight clothing.



3. Turn on side and keep airway clear.



4. Note the time a seizure starts and the length of time it lasts.



5. Don't put anything in mouth.



6. Don't hold down.

7. As seizure ends...offer help.

Step 3. Treatment options

Please record important information on the seizure log cards on the next page.

If I don't regain consciousness within _____ minutes, please:

- Call 911
- Once 911 is called, please call my emergency contacts below
 - If this box is checked advise EMTs that I have a VNS magnet

OR

Administer DIASTAT® AcuDial™ (diazepam rectal gel)

- For seizures that last more than _____ minutes
OR for _____ or more seizures in _____ hours
- My DIASTAT® AcuDial™ is kept: _____

Call 911 if

- I do not start waking up within _____ minutes after seizure is over (after giving DIASTAT® AcuDial™)
- Seizure does not stop within _____ minutes of giving DIASTAT® AcuDial™

Step 4. Notification

Call the following people if:

- I go to the Emergency Room
 You are concerned about my response
 DIASTAT® AcuDial™ is administered
 Other _____

Emergency Contacts

Name _____

Phone (____) _____

Name _____

Phone (____) _____

CONFIDENTIALITY STATEMENT

By creating and distributing this Pediatric Seizure Preparedness Plan, it is the expressed intention of Valeant Pharmaceuticals North America to provide a value-added service to healthcare providers, caregivers, and patients. Valeant Pharmaceuticals North America intends to fully comply with HIPPA guidelines for patient confidentiality to ensure that personal information is not disclosed to anyone outside of those that the parent/caregiver, healthcare professional, or student-patient deem acceptable. Valeant Pharmaceuticals North America is not responsible for this Plan or its contents after the Plan has been released to healthcare professionals. Careful consideration should be given before any medical information is released to others.

IMPORTANT SAFETY INFORMATION

DIASTAT® AcuDial™ (diazepam rectal gel) is a gel formulation of diazepam intended for rectal administration in the management of selected refractory patients with epilepsy, on stable regimens of AEDs, who require intermittent use of diazepam to control bouts of increased seizure activity for patients 2 years and older.

In clinical trials with DIASTAT®, the most frequent adverse event was somnolence (23%). Less frequent adverse events reported were dizziness, headache, pain, vasodilation, diarrhea, ataxia, euphoria, incoordination, asthma, rash, abdominal pain, nervousness and rhinitis (1%-5%). Please see full DIASTAT AcuDial Prescribing Information.

© 2007 Valeant Pharmaceuticals North America (INSERT PRC NUMBER: waiting on Client to provide)